

Analysis of Factors for Palliative Care for Breast Cancer Patients in Java in 2020

by Harnanik Nawangsari

Submission date: 16-Sep-2021 10:00AM (UTC+0700)

Submission ID: 1649577780

File name: ctors_for_Palliative_Care_for_Breast_Cancer_Patients_in_Java.pdf (153.4K)

Word count: 1822

Character count: 9129

Analysis of Factors for Palliative Care for Breast Cancer Patients in Java in 2020

Harmanik Nawangsari^{1,*}, Liliek Pratiwi²,

^{1,2} Lecturer of STIKES Insan Cendekia Medika Jombang, Lecturer of the Faculty of Health Sciences, Muhammadiyah University of Cirebon

*Corresponding author. Email: harnanik.nawangsari@gmail.com

ABSTRACT

This study pays attention to breast cancer patients in Java Province in 2020 because it aims to analyze factors related to palliative care, especially for patients with cervical cancer. So, it is hoped that this research will be useful as a suggestion for breast cancer patient health services. In this study, using a cross sectional research method. Research data was taken from several sources such as cancer foundations and family planning associations. This study uses a measuring instrument in the form of instruments from secondary data and a PNPC (Problem and Needs of Palliative Care) questionnaire. The sampling method in this study was purposive sampling, with a total sample of 178 female patients with breast cancer. Patients with breast cancer ≥ 1 year, age ≥ 20 years, able to write and read, in a conscious and stable state, were the inclusion criteria for this study. From the results of the study, with multivariate analysis, linear regression, the monthly income was obtained (with a value of $t = -2.90$, p value = 0.04, 95% CI = -3,789, -1,807), marital status ($t = -2.10$, p value = 0.04, 95% CI = -3,111, -1,665), place of domicile ($t = -5.10$, p value = 0.04, 95% CI = -4,211, -1,415), the patient also had difficulty accepting meaning or perceptions of death. Then breast cancer patients also felt difficult and confused to tell their complaints ($t = -6.10$, p value = 0.04, 95% CI = -6.211, -1.667, then the patient felt distressed and lacked knowledge about pain management ($t = -5.33$, p value = 0.03 95% CI = -3.211, 1.886. From these factors the results are related to the palliative care of breast cancer patients in Java, Indonesia in 2020, with an R value of 0.109. This, nurses, doctors, families must increase their role proactively. The motivation of the patient's family must also be studied in the future because the patient's family is actively participating by participating in the training of doctors and nurses who perform health services to these patients. For other researchers, it can also improve further study so that palliative care of women with breast cancer can run smoothly both at home and in the hospital.

Keywords: patients with breast cancer, Java province, 2020

1. INTRODUCTION

Every year and from time to time, there are additional cases of breast cancer, based on WHO data, there are 1.38 million new cases and 458 thousand cases of breast cancer that end in death. Women can be attacked by breast cancer and cervical cancer with various trigger factors. The impact of the development of breast cancer cells, decreasing the quality of life for patients and their families. There were 58,256 cases out of a total of 348,809 cancer cases. If there are 17 people per 1000 population affected by breast cancer, of course the impact must be considered. The number of deaths increases because it can be various aspects, whether it is the decreased quality of life of the patient, family motivation or quality of care. There is a known etiology of breast cancer, some need further study. 10% of the etiology of breast cancer, due to genetic consequences of the parents. The rest is in the range of 90% usually suddenly gets Covid19.

Excess of the hormone estrogen affects the growth of breast cancer cells in patients. In addition, individual factors are also related, such as how long the use of hormonal contraceptives has been and factors are not breastfeeding

and have never been pregnant. Previously and recently, from dietary factors, such as junk food in the long term and co-factors, can cause breast cancer. From the 2018 RISKESDAS data, cancer and tumors increased from 2013. Indonesia as a state in Southeast Asia is a country with 8th and 23rd rank in Asia, as a country with a number of cancer cases. Breast cancer is 42.1 per 100,000 population and is a concern of the public and medical personnel.

The cure rate for breast cancer patients varies depending on the severity and whether there are comorbidities. Research on the motivation, knowledge, attitudes of breast cancer patients has been done a lot, but novelty, but not yet studied the extent to which palliative care is carried out and its factors. Patients need partner in maintaining their health. In some consumer accounts the patient's family is here.

2. METHOD

PNPC-sv consists of 33 question items related to problems and in reducing bias in this study, the number of samples in this study was taken 178 so that it could represent women with breast cancer in Java in 2020. The instrument in this study, namely the PNPC is a self-

assessment. with seven domains, namely daily activities, physical symptoms, autonomy, social issues, psychological issues, spiritual issues and financial issues, the need for palliative care. Each item will be asked about the problem and from the perceived problem whether they need help from health workers with the answer choices are (yes, many / no). PNP has a significant correlation with the dimensions of quality of life developed by EORTC QLQ-C30 (European Organization for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30) with p value = 0.01. PNP also has good internal consistency Alpha value of 0.70-0.86. PNP also has good internal consistency.

3. RESULT

Based on the results of multivariate analysis with linear regression, it was found that monthly income ($t = -2.90$, p value = 0.04, 95% CI = -3.789- 1.807), marital status ($t = -2.10$, p value = 0.04, 95% CI = -3,111- 1,665), place of residence ($t = -5.10$, p value = 0.04, 95% CI = -4,211- 1,415), difficulty perceiving the meaning of death ($t = -6.10$, p value = 0.07, 95% CI = -9,211- 1,110), difficulty finding someone to tell stories or talk to ($t = -6.10$, p value = 0.04, 95% CI = -6,211- 1,677) difficulty in pain management ($t = -5.33$, p value = 0.03, 95% CI = -3.211- 1.886) associated with palliative care for women with breast cancer in Indonesia in 2020, the R value is 0.109. In this study, it was found that the overall needs of cancer patients were in the upper middle class. From the results of the study, with multivariate analysis, linear regression, the monthly income was obtained (with a value of $t = -2.90$, p value = 0.04, 95% CI = -3,789, -1,807), marital status ($t = -2.10$, p value = 0.04, 95% CI = -3,111, -1,665), place of domicile ($t = -5.10$, p value = 0.04, 95% CI = -4,211, -1,415), the patient also had difficulty accepting meaning or perceptions of death. Then breast cancer patients also felt difficult and confused to tell their complaints ($t = -6.10$, p value = 0.04, 95% CI = -6.211, -1.667, then the patient felt distressed and lacked knowledge about pain management ($t = -5.33$, p value = 0.03 95% CI = -3.211, 1.886. From these factors the results are related to the palliative care of breast cancer patients in Java, Indonesia in 2020, with an R value of 0.109., 2015 that breast cancer patients hope that health workers who care for them can honestly say their condition and follow up their needs that have not been met. dimensions, as well as patient-oriented care (Alikhani, 2019) In De Silva M's (2010) study, women who previously had a history of breastfeeding were healthier and less prone to breast cancer. From the research results, it was also stated that the use of health care facilities also affects the palliative needs of breast cancer patients. Society and community can play a more dominant role in hospital services for breast cancer patients. So, public awareness about the importance of palliative care must be raised again.

4. CONCLUSION

From the results of this study, it is better if the questionnaire regarding palliative care can be socialized to the hospital.

So, the hospital can improve the quality of health services. So, the advice for health service providers is not only medical personnel, but also families and communities in the form of comprehensive palliative service motivation.

ACKNOWLEDGMENT

The researchers would like to thank all those who have helped the course of this research until the research was published.

REFERENCES

- [1] Anindita Mukherjee, Koustav Mazumder, Sushmita Ghoshal. Impact of different sociodemographic factors on mental health status of female cancer patients receiving chemotherapy for recurrent disease. 2018. 24 (4). (426-440). Indian Journal of Palliative Care.
- [2] Beasley JM, Kwan ML, Chen WY, et al. Meeting the physical activity guidelines and survival after breast cancer: findings from the after breast cancer pooling project. *Breast Cancer Res Treat.* 2012; 131: 637- 643.
- [3] Bradshaw PT, Ibrahim JG, Khankari N, et al. Post-diagnosis physical activity and survival after breast cancer diagnosis: The Long Island Breast Cancer Study. *Breast Cancer Res Treat.* 2014; 145: 735- 742.
- [4] Loprinzi PD, Cardinal BJ, Winters-Stone K, et al. Physical activity and the risk of breast cancer recurrence: a literature review. *Oncol Nurs Pract.* 2012; 39: 269- 274.
- [5] Darby SC, Ewertz M, McGale P, Bennet AM, Blom-Goldman U, Bronnum D, et al. Risk of ischemic heart disease in women after radiotherapy for breast cancer. *N Engl J Med.* 2013;368(11):987e98. <http://dx.doi.org/10.1056/NEJMoa1209825>
- [6] World Health Organization. Breast cancer: prevention and control [cited 2012 June 3]. Available from: <http://www.who.int/cancer/detection/breastcancer/en/index.html>
- [7] Morrow M, Burstein HJ, and Harris JR. Malignant tumors of the breast. De Vita VT Jr, Lawrence TS, & Rosenberg SA. (2015). *Cancer: Principles and Practice of Oncology.* (10th Edition). Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins. 79: 1117-1156
- [8] National Toxicology Program. (2016). *14th Report on Carcinogens.* Department of Health and Human Services. Retrieved from: <http://ntp.niehs.nih.gov/pubhealth/roc/index-1.html>.
- [9] Wentlandt K, Krzyzanowska MK, Swami N, Rodin GM, Le LW, Zimmermann C. Referral practices of oncologists to specialized palliative care. *J Clin Oncol* 2012;30: 438-45.
- [10] Hui D, Kim YJ, Park JC, Zhang Y, Strasser F, Cherny N, et al. Integration of oncology and palliative care: a thematic review. *Oncologist* 2015;20:77-83.
- [11] Oesman I. Dampak Kanker Payudara. In: Indah RSP, editor. Jakarta: PT Meta Publishing; 2012.
- [12] De Silva M, Senarath U, Gunatilake M, Lokuhetty D. Prolonged breastfeeding reduces risk of breast cancer in Sri Lanka women. *Cancer Epidemiol.* 2010;34(3):267-73.
- [13] Mahtab Alikhani, Soudabeh Vatankhah, Hasan Abolghasem Gorji, Hamid Ravaghi. A Comparison of Policy Analysis of Palliative Care for Cancer in UK, Malaysia, and South Africa. *Int J Cancer Manag.* 2019. 12(12):e94841

Analysis of Factors for Palliative Care for Breast Cancer Patients in Java in 2020

ORIGINALITY REPORT

18%

SIMILARITY INDEX

17%

INTERNET SOURCES

15%

PUBLICATIONS

15%

STUDENT PAPERS

PRIMARY SOURCES

1	Ping-Ping Bao, Gen-Ming Zhao, Xiao-Ou Shu, Peng Peng, Hui Cai, Wei Lu, Ying Zheng. "Modifiable Lifestyle Factors and Triple-negative Breast Cancer Survival", <i>Epidemiology</i> , 2015 Publication	2%
2	bmccancer.biomedcentral.com Internet Source	2%
3	bjsm.bmj.com Internet Source	2%
4	Submitted to University of College Cork Student Paper	2%
5	Submitted to University of Arizona Student Paper	2%
6	www.iosrjournals.org Internet Source	1%
7	link.springer.com Internet Source	1%

8	Submitted to University of Wisconsin, La Crosse Student Paper	1 %
9	dergipark.org.tr Internet Source	1 %
10	Submitted to St. Mary's College Twickenham Student Paper	1 %
11	Submitted to University of Central Florida Student Paper	1 %
12	scholar.unand.ac.id Internet Source	1 %
13	www.thieme-connect.com Internet Source	<1 %
14	bmcpalliatcare.biomedcentral.com Internet Source	<1 %
15	Sato, Kazuhiko, Yoshio Mizuno, Hiromi Fuchikami, Masahiro Kato, Takahiro Shimo, Jun Kubota, Naoko Takeda, Yuko Inoue, Hiroshi Seto, and Tomohiko Okawa. "Individualized Case–Control Study of Accelerated Partial Breast Irradiation by Multicatheter Intertstitial Brachytherapy in Japanese Patients with Breast Cancer", Cancer and Clinical Oncology, 2013. Publication	<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off